

## SPHERO™ Coated Magnetic Particles

- Available with a variety of ligands such as Streptavidin, Avidin, Neutravidin, Protein A, Protein G, and Biotin
- Also available coated with highly specific recognition groups such as polyclonal antibodies
- Used in nucleic acid isolation, protein purification, immunology, and cell separations
- Available impregnated with fluorophores for flow cytometry or easy particle location identification in phagocytosis assays.

Magnetic particles coated with Avidin, Streptavidin, Biotin, Protein A and various antibodies are available from Spherotech. All of the proteins used are covalently coupled to the magnetic particles. The coated magnetic particles are supplied as a suspension in phosphate buffer, pH 7.4 with 0.02% sodium azide (some products also contain 0.1% BSA). Please refer to the recommended coating procedures on page 88 for more detailed technical information and coating.

Similarly to the magnetic particles offered on page 61 Spherotech coated magnetic particles are offered as the classic, encapsulated, or crosslinked magnetic microsphere. See pages 62 for benefits of each type.

## SPHERO™ Biotin Coated Magnetic Particles

- Used to take advantage of the high affinities of the biotin–streptavidin and biotin–avidin interactions ( $K_a$  in the order of  $10^{13}$ – $10^{15}$   $M^{-1}$ )
- Represents one of the strongest biomolecules interactions to form stable complexes.

Particle Type and Surface	Size, $\mu m$	% w/v	Catalog No.	Unit
Biotin	1.0-1.4	0.5	TM-10-10	10 mL
Biotin	4.0-4.5	1.0	TM-40-10	10 mL
Biotin	6.0-7.9	1.0	TM-60-5	5 mL
Biotin, Cross-linked, granules, non-uniform	~1-2 $\mu m$	0.5	TMX-10-10	10 mL

## SPHERO™ Avidin Coated Magnetic Particles

- Used for Genome isolation when coated with a biotinylated genome capture probe for E.coli and B.subtilis\*  
\*S.Yeung, T. Ming-Hung Lee, H. Cai, and I-Ming Hsing. "A DNA biochip for on-the-spot multiplexed pathogen identification." Nucleic Acids Res., Vol 34, No. 18, e118 (Oct 2006)
- See pages 70 and 71 for uses of streptavidin and avidin coated particles.

Particle Type and Surface	Size, $\mu m$	% w/v	Catalog No.	Unit
Avidin	1.0-1.4	0.5	VM-10-10	10 mL
Avidin	4.0-4.5	1.0	VM-40-10	10 mL
Avidin	4.0-4.5	1.0	VM-40-100	100 mL
Avidin	6.0-8.0	1.0	VM-60-10	10 mL
Avidin	6.0-8.0	1.0	VM-60-100	100 mL
Avidin	8.1-9.9	1.0	VM-80-5	5 mL
Avidin, Smooth Surface	3.0-3.9	1.0	VMS-30-10	10 mL
Avidin, Smooth Surface	4.0-5.0	1.0	VMS-40-10	10 mL
Avidin, Cross-linked, granules, non-uniform	~1-2 $\mu m$	0.5	VMX-10-10	10 mL
Avidin, Cross-linked, granules, non-uniform	~1-2 $\mu m$	0.5	VMX-10-100	100 mL

## SPHERO™ Streptavidin Coated Magnetic Particles

- Streptavidin magnetic particles have found widespread use as detection reagents in immunology, biochemistry and cell biology due to their high affinity binding to biotin
- Biotin–streptavidin interaction have been exploited in many applications including the development of new reagents for diagnostics such as sandwich magnetic particle enzyme-linked immunosorbent assay (MPEIA) and molecular biology studies involving nucleic acids.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Streptavidin, High Iron	0.2-0.39	0.5	SVM-025-5H	5 mL
Streptavidin	0.4-0.69	0.5	SVM-05-10	10 mL
Streptavidin, High Iron	0.4-0.69	0.5	SVM-05-5H	5 mL
Streptavidin	0.7-0.9	0.5	SVM-08-10	10 mL
Streptavidin	1.0-1.4	0.5	SVM-10-10	10 mL
Streptavidin	1.5-1.9	0.5	SVM-15-10	10 mL
Streptavidin	2.0-2.9	0.5	SVM-20-10	10 mL
Streptavidin	3.0-3.9	1.0	SVM-30-10	10 mL
Streptavidin	4.0-4.5	1.0	SVM-40-10	10 mL
Streptavidin	4.6-5.9	1.0	SVM-50-5	5 mL
Streptavidin	6.0-7.9	1.0	SVM-60-5	5 mL
Streptavidin	8.0-9.9	1.0	SVM-80-5	5 mL
Streptavidin, High Iron	38.0-44.0	0.5	SVMH-400-4	4 mL
Streptavidin, High Iron	45.0-52.0	0.5	SVMH-500-4	4 mL
Streptavidin, Smooth Surface	3.0-3.9	1.0	SVMS-30-10	10 mL
Streptavidin, Smooth Surface	4.0-5.0	1.0	SVMS-40-10	10 mL
Streptavidin, Cross-linked, granules, non-uniform	~1-2 $\mu\text{m}$	0.5	SVMX-10-10	10 mL

## SPHERO™ Neutravidin Coated Magnetic Particles

- Provide very low non-specific binding since they do not contain any carbohydrates and have a near-neutral isoelectric point of 6.3
- Used for diagnostic and molecular biology applications
- Have a binding capacity of ~5100 pmol/mg.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Neutravidin	2.0-2.9	1.0	NVM-20-5	5 mL

## SPHERO™ Rabbit anti-HA Coated Magnetic Particles

- Coated with affinity purified anti-Hemagglutinin (HA) epitope tag [Rabbit] polyclonal antibody
- Binds to HA-tagged recombinant proteins.

Particle Type and Surface	Size, $\mu\text{m}$	Catalog No.	Unit
Rabbit anti-HA, Smooth Surface, $10^7/\text{mL}$	3.0-3.9	RHAMS-30-2	2 mL

**SPHERO™ Protein G Coated Magnetic Particles**

- Used to link capture species-specific anti-IgG to magnetic microspheres
- Directly binds immunoglobulins from ascites fluids or concentrated hybridoma supernatants to facilitate purification.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Protein G	4.0-4.5	1.0	PGM-40-5	5 mL
Protein G, Smooth Surface	3.0-3.9	1.0	PGMS-30-5	5 mL
Protein G, Smooth Surface	4.0-5.0	1.0	PGMS-40-5	5 mL
Protein G, Cross-linked, granules, non-uniform	~1-2	1.0	PGMX-10-5	5 mL

**SPHERO™ Protein A Coated Magnetic Particles**

- Used for the immunomagnetic separation (IMS) and real-time PCR to detect *Escherichia coli*\*

\*Fu, Z., S. Rogelj, et al. (2005). "Rapid detection of *Escherichia coli* O157:H7 by immunomagnetic separation and real-time PCR." *International Journal of Food Microbiology* 99(1): 47-57.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Protein A	4.0-4.5	1.0	PAM-40-5	5 mL
Protein A, Smooth Surface	3.0-3.9	1.0	PAMS-30-5	5 mL
Protein A, Smooth Surface	4.0-5.0	1.0	PAMS-40-5	5 mL
Protein A, Cross-linked, granules, non-uniform	~1-2	1.0	PAMX-10-5	5 mL

**SPHERO™ Sheep anti-Rat IgG Coated Magnetic Particles**

- Consists of uniform, paramagnetic polystyrene beads coated with polyclonal Sheep anti-Rat IgG antibodies.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Sheep anti-Rat IgG (H&L)	4.0-4.5	2.0	SRM-40-5	5 mL

**SPHERO™ Donkey anti-Goat IgG Coated Magnetic Particles**

- Ideal for direct or indirect isolation of proteins and cells during immunomagnetic separation.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Donkey anti-Goat IgG (H&L) Cross-adsorbed	4.0-4.5	1.0	GMXA-40-10	10 mL


**SPHERO™ Goat anti-Rabbit IgG Coated Magnetic Particles**

- Used in immunomagnetic separation (IMS)\*

\*Antognoli, M. C., M. D. Salman, et al. (2001). "A one-tube nested polymerase chain reaction for the detection of *Mycobacterium bovis* in spiked milk samples: an evaluation of concentration and lytic techniques." *J Vet Diagn Invest* 13(2): 111-116.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Goat anti-Rabbit IgG (H&L)	4.0-4.9	1.0	RM-40-10	10 mL
Goat anti-Rabbit IgG (Fc)	4.0-4.5	1.0	RMFc-40-10	10 mL
Goat anti-Rabbit IgG (H&L), Smooth Surface	3.0-3.9	1.0	RMS-30-10	10 mL
Goat anti-Rabbit IgG (H&L), Smooth Surface	4.0-5.0	1.0	RMS-40-10	10 mL
Goat anti-Rabbit IgG (Fc), Smooth Surface	3.0-3.9	1.0	RMSFc-30-10	10 mL
Goat anti-Rabbit IgG (H&L), Cross-linked, granules, non-uniform	~1-2 $\mu\text{m}$	0.5	RMX-10-10	10 mL

**Figure 113** Size distribution analysis of SPHERO™ Cat. No. RM-40-10, Gt anti-Rb IgG (H&L) Magnetic Particles from the Beckman Coulter Z3 Multisizer™.



**SPHERO™ Goat anti-Mouse IgG Coated Magnetic Particles****Attributes**

- Uniform particle size
- Paramagnetic in nature
- Rapid magnetic responsiveness
- Low non-specific binding
- High binding capacity
- Consistent lot-to-lot performance.

**Applications**

- Automated immunoassays
- Immunoprecipitation
- IP-western blots.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Goat anti-Mouse IgG (H&L)	4.0-4.5	1.0	MM-40-10	10 mL
Goat anti-Mouse IgG (Fc)	4.0-4.5	1.0	MMFc-40-10	10 mL
Goat anti-Mouse IgG (H&L), Cross adsorbed	4.0-4.5	1.0	MMXA-40-10	10 mL
Goat anti-Mouse IgG (H&L), Smooth Surface	3.0-3.9	1.0	MMS-30-10	10 mL
Goat anti-Mouse IgG (H&L), Smooth Surface	4.0-5.0	1.0	MMS-40-10	10 mL
Goat anti-Mouse IgG (Fc), Smooth Surface	3.0-3.9	1.0	MMSFc-30-10	10 mL
Goat anti-Mouse IgG (Fc), Smooth Surface	4.0-5.0	1.0	MMSFc-40-10	10 mL
Goat anti-Mouse IgG (H&L), Smooth Surface, Cross adsorbed	3.0-3.9	1.0	MMSXA-30-10	10 mL
Goat anti-Mouse IgG (H&L), Smooth Surface, Cross adsorbed	4.0-5.0	1.0	MMSXA-40-10	10 mL
Goat anti-Mouse IgG (H&L), Cross-linked, granules, non-uniform	~1-2 $\mu\text{m}$	0.5	MMX-10-10	10 mL
Goat anti-Mouse IgG (H&L), Cross adsorbed, Cross-linked, granules, non-uniform	~1-2 $\mu\text{m}$	0.5	MMXA-10-10	10 mL

**SPHERO™ Goat anti-Human IgG Coated Magnetic Particles**

- Ideal for the capture and/or detection of target analytes by direct or indirect isolation during immunomagnetic separation
- Improves the performance of ELISAs by enhancing sensitivity and shortening incubation times.

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Goat anti-Human IgG (H&L)	4.0-4.5	1.0	HM-40-10	10 mL
Goat anti-Human IgG (H&L), Smooth Surface	3.0-3.9	1.0	HMS-30-10	10 mL
Goat anti-Human IgG (H&L), Smooth Surface	4.0-5.0	1.0	HMS-40-10	10 mL
Goat anti-Human IgG (H&L), Cross-linked, granules, non-uniform	~1-2 $\mu\text{m}$	0.5	HMX-10-10	10 mL

**SPHERO™ Con A Coated Magnetic Particles**

- Binds to saccharide functional groups on the cell surface\*

Particle Type and Surface	Size, $\mu\text{m}$	% w/v	Catalog No.	Unit
Con A	0.7-0.9	1.0	CAM-08-10	10 mL

\*Gupta, S., R. G. Alargova, et al. "On-Chip Dielectrophoretic Coassembly of Live Cells and Particles into Responsive Biomaterials." *Langmuir*, 2010, 26 (5), pp 3441–3452.